

ISSN 1392-0146 (print)  
ISSN 2029-0578 (online)  
www.lmaleidykla.lt



# Biologija

---

*Bendroji, molekulinė ir ląstelės biologija • Biochemija • Biofizika •  
Biotechnologija • Bendroji, molekulinė ir landšaftų ekologija •  
Antropogeninis poveikis ekosistemoms • Ekosistemų apsauga •  
Genetika • Mikrobiologija • Imunologija •  
Parazitologija • Fiziologija •*

---

*General, Molecular and Cell Biology • Biochemistry • Biophysics •  
Biotechnology • General, Molecular and Landscape Ecology •  
Anthropogenic Impact on Ecosystems • Ecosystem Protection •  
Genetics • Microbiology • Immunology •  
Parasitology • Physiology •*

---

**2020 Vol. 66 No. 1**



## Editorial Board | Redakcinė kolegija

### Editor-in-Chief | Vyriausiasis redaktorius

**Algimantas PAULAUSKAS**

Faculty of Natural Sciences

Vytautas Magnus University

Vileikos St. 8, 44404 Kaunas, Lithuania

Email: biologija@vdu.lt

### Vice-editor | Vyriausiojo redaktoriaus pavaduotojas

**Vincas BŪDA**

Nature Research Centre, Lithuania

(ecology, entomology)

### Editorial Assistant | Atsakingoji sekretorė

**Indrė LIPATOVA**

Vytautas Magnus University

Vileikos St. 8, 44404 Kaunas, Lithuania

Email: biologija@vdu.lt

Tel: +370 37 327905, Fax: +370 37 327916

## Editorial Board | Redakcinė valdyba

Aleksandr ALIMOV, Zoological Institute of the Russian Academy of Sciences, Russia (*biodiversity and ecology of hydrobionts*)

Makoto ASASHIMA, University of Tokyo, Japan (*developmental biology, cell biology, organogenesis*)

Anna-Karin BORG-KARLSON, Royal Institute of Technology, Sweden (*molecular biology, biochemistry*)

Vilmantė BORUTAITĖ, Lithuanian University of Health Sciences, Lithuania (*biochemistry*)

Vladas Algirdas BUMELIS, State Research Institute Centre for Innovative Medicine, Lithuania (*biotechnology*)

Dalius BUTKAUSKAS, Nature Research Centre, Vilnius, Lithuania (*molecular ecology*)

Michael A. FULLEN, University of Wolverhampton, Central England, UK (*soil management, ecology*)

Yuri Y. GLEBA, Icon Genetics GmbH, Halle/Saale, Germany (*molecular biology, plant biotechnology, gene engineering*)

Rienk van GRONDELLE, Free University of Amsterdam (The Vrije Universiteit Amsterdam), Netherlands, (*biophysics*)

Stephen E. HALFORD, University of Bristol, United Kingdom, (*nucleic acids, enzymes, biochemistry*)

Sergei G. INGE-VECHTOMOV, St. Petersburg State University, Russia (*microbiology, molecular biology, genetics*)

Robert Neil JONES, University of Wales, Aberystwyth, United Kingdom (*plant molecular biology, cytogenetics*)

Victor V. KLEMAS, University of Delaware, USA (*ecosystem protection*)

Skirmantas KRIAUCIONIS, Ludwig Institute for Cancer Research Ltd, University of Oxford, United Kingdom (*animal biotechnology, gene engineering, genetics*)

Juozas KULYS, Vilnius Gediminas Technical University, Lithuania (*molecular biology, biochemistry, biotechnology, bioinformatics*)

Vladimir V. KUZNETSOV, Russian Academy of Sciences, Timiryazev Institute of Plant Physiology, Russia (*plant physiology, biochemistry*)

Juozas Rimantas LAZUTKA, Vilnius University, Lithuania (*genetics, general biology*)

Ivan Karl-Eric MAGNUSSON, Linköping University, Sweden (*cell biology, biochemistry, immunology*)

Jonas Remigijus NAUJALIS, Vilnius University, Lithuania (*plant biology and ecology*)

Sten NILSSON, International Institute for Applied Systems Analysis (IIASA), Austria (*anthropogenic impact on ecosystem*)

Philippos C. PATSALIS, Cyprus Institute of Neurology and Genetics, Cyprus (*neurology, cytogenetics*)

Dmitrij PAVLOV, A. N. Severtsov Institute of Ecology and Evolution of the Russian Academy of Sciences, Russia (*ichthyology, fish ecology*)

Branislav PEŤKO, Parasitological Institute of the Slovak Academy of Sciences, Slovakia (*ecology, entomology*)

Isaaks RASHAL, Institute of Biology, University of Latvia (*plant biology, genetics*)

Vladimir N. RESHETNIKOV, National Academy of Sciences of Belarus, Central Botanical Garden (*plant physiology, biochemistry, biotechnology, immunology*)

Karl SKIRNISSON, University of Iceland, Iceland (*zoology, parasitology*)

Eva Maria SEVERINSON, Stockholm University, Sweden (*immunology*)

Gediminas VALKIŪNAS, Nature Research Centre, Lithuania (*zoology, parasitology*)

Alexei VERKHRATSKY, University of Manchester, United Kingdom, (*cellular neurophysiology*)

Richard VILLEMS, University of Tartu, Estonia (*genetics, general biology*)

Vytas Antanas TAMOŠIŪNAS, State Research Institute Centre for Innovative Medicine, Lithuania (*immunology, cell biology*)

**Published:** 2017-09-04

## General Biology

- [\*\*The mystery of the tiny Urticaceae-feeders: documentation of the first leaf-mining Nepticulidae \(Lepidoptera\) species from equatorial America associated with Phenax, Boehmeria and Pilea\*\*](#)

Jonas R. Stonis, Arūnas Diškus, Andrius Remeikis, Alex K. Monro

○ [PDF](#)

- [\*\*A study of epidermal leaf anatomy of 18 Euphorbia taxa from Kerman Province, Iran\*\*](#)

Seyed Mehdi Talebi, Mitra Noori, Habibeh Afzali Naniz

○ [PDF](#)

## Ecology

- [\*\*Factors influencing seed germination of the pastoral plant Retama raetam subsp. bovei \(Fabaceae\): interactive effects of fruit morphology, salinity, and osmotic stress\*\*](#)

Kaouther Mechergui, Hela Mahmoudi, Mohamed Larbi Khouja, Wahbi Jaouadi

○ [PDF](#)

## Molecular Biology

- [\*\*Effect of alimentary synthetic estrogen on cell compensatory mechanisms in rats of different ages\*\*](#)

Oleh Yermishev, Tatyana Lykholat, Olena Lykholat

○ [PDF](#)

## Microbiology

- [\*\*Effect of amino acids on pigments, citrinin, and lovastatin production by Monascus purpureus under static conditions\*\*](#)

Padmavathi Talapragada, Rashmi Dikshit, Mudaheranwa Phocas, Madhusudan. M.R., Sanjana Samprathi

○ [PDF](#)

## Parasitology

- [\*\*Laelapidae mites \(Acari: Mesostigmata\) infesting small rodents in the Curonian Spit, Lithuania\*\*](#)

Evelina Kaminskienė, Jana Radzijeuskaja, Linas Balčiauskas, Vaclovas Gedminas, Algimantas Paulauskas

○ [PDF](#)

- [\*\*The effect of thyme \(\*Thymus caucasicus\*\) ethanol extract on Varroa mite \(\*Varroa destructor\*\), an ectoparasite mite of \*Apis mellifera meda\* \(Hym: Apidae\)\*\*](#)

Ataollah Rahimi, Yaser Khoram Del, Farzad Moradpour

○ [PDF](#)

Physiology

- [\*\*Activity and isoenzyme composition of peroxidase in the vegetative organs of Japanese quince under steppe zone conditions\*\*](#)

Yuriy Prisedsky, Anatoliy Kabar, Yuriy Lykholat, Nadezhda Martynova, Larysa Shupranova

○ [PDF](#)

- [\*\*Climate warming caused changes in the bud swelling phenophase of introduced conifer species, considering the aspect of hardiness zones in their natural habitats\*\*](#)

Asta Malakauskienė, Adomas Vitas

○ [PDF](#)

Biophysics

- [\*\*Influence of NMDA and GABA synaptic dysfunction on the evoked gamma oscillations in a computational model of schizophrenia\*\*](#)

Rokas Jackevičius, Bruce P. Graham, Aušra Saudargienė

○ [PDF](#)

Biotechnology

- [\*\*Design of a chitosan-based bio-scaffolding for regenerative medicine: compatibility analysis for cartilage regeneration\*\*](#)

Seyed Ali (Behruz) Khaghani, Gunay Akbarova, Gulrukh Dilbazi

○ [PDF](#)

# Effect of alimentary synthetic estrogen on cell compensatory mechanisms in rats of different ages

Oleh Yermishev<sup>1</sup>,

Tatyana Lykholat<sup>2</sup>,

Olena Lykholat<sup>2\*</sup>

<sup>1</sup> *Vasyl Stus Donetsk National University,  
600-richya St. 21,  
21021 Vinnitsa, Ukraine*

<sup>2</sup> *Oles Honchar Dnipropetrovsk  
National University, Haharina Ave. 72,  
49000 Dnipropetrovsk, Ukraine*

Results of a study of lipid peroxidation, antioxidant system components, and cholinergic neurotransmitter system in the organs of experimental rats of different ages exposed to alimentary synthetic estrogen are presented. Given the state of peroxidation processes and AChE activities in female rats exposed to xenoestrogen, it is possible to assume the possibility of the restructuring of the functioning of mediator and enzyme systems and additional strengthening of pathological symptoms. In the future, such phenomena may trigger the reduction of potential of compensatory mechanisms in compromising the health of the consumers. In puberty, females were more sensitive to nutritional synthetic estrogen than mature animals, thus proving that age is another factor in xenoestrogen exposure. Because of the changes in the rates of reactions to detoxification but not to the metabolism of estrogen received into the organism, particularly with food, with age the animals were less susceptible to the effects of these substances.

**Keywords:** xenoestrogen, lipid peroxidation, antioxidant system, cholinergic regulation, age

## INTRODUCTION

Many contaminants are associated with their potential estrogenic effect and classified as endocrine-disrupting compounds, including numerous classes of organic compounds. Substances-depleted endocrine systems are a compound changing the hormonal and homeostatic system and acting through different mechanisms (Darbre, Charles, 2010).

Estrogens are female sex hormones produced by the follicles of the ovaries, placenta, partially adrenal cortex, and the testes. They regulate specific sexual functions. Estrogens spread through

the bloodstream to various tissues, including the cardiovascular, the immune, and the central nervous systems. Due to their lipophilic character they can easily diffuse through cell membranes and hematoencephalic barrier exerting effects (Petrone et al., 2014). Estrogens may lead to structural changes in the body, including a modulation of mitochondrial ATP synthesis and of the scope of protein biosynthesis (Shearwood et al., 2015). The activity of steroid hormones is targeted at the brain. They can influence the architecture and plasticity of the brain, the neurochemical and molecular processes, as well as the behaviour and neural protection through genomic and non-genomic action. These processes are important for brain activity (McEwen, 2014).

\*Corresponding author. E-mail: lykholat2010@ukr.net

Xenoestrogens are industrial chemicals (for example, esters of phthalic acid, bisphenol A, phenolic compounds, and parabens) used in the manufacture of polycarbonate plastics, epoxy resins, food packaging and coatings, dental sealants, rubber chemicals, and flame retardants (Arditsoglou, Voutsas, 2008; Meulenberg, 2009; Boberg et al., 2010; Rochester, 2013). Similar compounds are present in antimicrobial substances (zearalenone) and medicines, personal hygiene products (Wielogórska et al., 2015), drinking and bottled water (Dhaini, Nassif, 2014), foods – grains, fruits and vegetables (Massart, Saggese, 2010), animal products – meat, dairy, eggs (Kolf-Clauw et al., 2008; Wielogorska et al., 2014). A special group is food containing phytoestrogens (Tomar, Shiao, 2008). Xenobiotic substances are contained in food (inhibitors of proteolytic enzymes) originally or through contamination with herbicides, pesticides, animal growth regulators, etc., some of which can form during improper cooking (roasting or smoking) (Hulin et al., 2014). Despite strict EU laws on the need of ensuring the safety of supply in view of low concentration levels of residual pesticides, the threat of pesticide residues exists in baby food (Hercegová et al., 2007). Such nutritional factors as phytoestrogens are contained in food, and legumes are particularly rich in them (Cassidy, 2003).

Consumers are put at the risk of low levels of exposure to these substances throughout their lives. At least half of these substances are resistant to natural degradation. The effect of these compounds is being evaluated with a focus on mutagenicity and genotoxicity. However, this approach ignores the recent integration of new toxicological studies such as endocrine disorders, mixture toxicity, and toxicity to development. Although most of the environmental estrogen activity was recognized as “weak” for many years because of its inability to cause transcriptional effects, at present it has been proven that they are quite powerful initiators of signal cascades out of membranes (De Coster, van Larebeke, 2012).

The environmental hazard of these compounds lies in their ability to disrupt or modify the endocrine system of living organisms, which

is central to their functioning, development and reproduction. They interact with a normally functioning endocrine system, modulate or block the action of natural hormones in the body, launch reactions identical to them thus affecting the synthesis, secretion, transport, binding, action, or elimination of natural hormones. They are active in very low doses, especially in foetuses and newborns. Endocrine-disrupting substances contribute significantly to the occurrence of cancer, neurological and behavioral disorders, the number of which has increased markedly in recent decades.

Hormones have different effects on the body at the different periods of life and this is due not only to age hormonal homeostasis, but also to the state of hormone-dependent organs (Hu et al., 2013; Hulin et al., 2014). The onset of puberty is characterized by a significant increase in the synthesis of estrogen by the ovaries due to changes in the “settings” of the hypothalamus, which “allows” a high concentration of the hormone in the blood.

Lipid peroxidation (LPO) is a universal biological mechanism constantly occurring in cellular membranes. Free radicals are mediators of signal transduction as a component of fundamental mechanism of inflammation, cytokines induction, proliferation, apoptosis, gene regulation of cellular security, chromatin remodulation, and poly(ADP-ribose) synthase activation. The recurrent peroxidation syndrome is a factor of the pathogenesis of several diseases: it serves as the basis for allocating them to the free radical pathology. The main components of the enzymatic antioxidation systems are superoxide dismutase (SOD) inactivating oxygen radicals, catalase, and glutathione redox system. Violation of the sensitive balance between free radicals and antioxidants can result in cell damage and trigger disorders of homeostasis (Lykholat et al., 2016). Exposure of xenoestrogen leads to strengthening of the generation of hydroxyl radicals with ensuing histopathological changes in the liver and kidneys, which can be explained either by estrogenic activity or the inclusion of different ways causing oxidative stress (Rashid et al., 2009).

Acetylcholinesterase (AChE) plays a key role in the neurohumoral and synaptic transmission: it catalyzes the hydrolysis of acetylcholine in the cholinergic synapses and thus terminates this neurotransmitter act on cholinergic receptors responsible for the permeability of the postsynaptic membrane for ions. The system acetylcholine-acetylcholinesterase is one of the main parts of the autonomic nervous system that involves other mediator and modulator processes performing the reactions of effector organs on central nervous impulses.

Risks associated with nutritional influence of synthetic hormones have not been fully characterized yet, and publications give conflicting reports about long-term consequences of the use of such products (Omoruyi et al., 2013). Given the threat of contamination by chemicals with endocrine properties in the diet of pubertal children as a particularly sensitive population, an analysis of the risk regarding the occurrence of health problems in the future is required. The potential estradiol effects unrelated to reproductive functions require further study (Courant et al., 2008; Kummer et al., 2012; Hu et al., 2013; Hulin et al., 2014). The membrane acetylcholinesterase activity, cell stability, the anti-oxidative activity, and the generation of free radicals are the parameters used to characterize structural and functional changes in cells (Kujawa et al., 2014).

**The aim** of the present work was the study lipid peroxidation, antioxidant system components, and cholinergic neurotransmitter system in the organs of experimental rats of different ages for exposure to alimentary synthetic estrogen.

## MATERIALS AND METHODS

The experiments were conducted on Wistar rats exposed to exogenous estrogen for 45 days. At the beginning of the experiment, three-month-old pubertal animals (group 2) and six-month-old sexually mature rats (group 4) were involved. The control group consisted of intact animals of corresponding age (groups 1 and 3).

For the modelling of exogenous estrogen impact, the rats' food was treated with the drug

Sinestrol as stilbene derivative that differs from estrogen steroid hormones in chemical structure, but is similar to them in biological and medicinal properties at the rate of 2 mg/kg.

Research materials were the brain, the liver, kidneys, and blood serum of rats. The objects were indices of TBA-active products, the total antioxidant activity, the activity of superoxide dismutase, glutathione peroxidase, glutathione transferase, glutathione reductase, glutathione content of the recovery, and AchE activity on standard methods.

The data were treated with standard methods of variational series estimation. The difference between the comparative values was considered significant at  $p < 0.05$ .

## RESULTS

The level of TBA-active products in the liver exceeded the control index by 26% in the group of pubertal rats and by 18.5% in mature females. In kidneys, the accumulation of TBA-active products exceeded the control index by 10% (group 2) and by 21.5% (group 4). Activity of lipid peroxidation in the brain increased by 34% in experimental group 2, and by 29.5% in group 4. The TBKAP level exceeded the control index by 48% in the serum of 4.5-month-old rats and by 51% in the serum of 7.5-month-old rats (Table 1).

Glutathione is one of the most powerful cellular antioxidants. Glutathione deficiency is observed in numerous diseases including tumours. The mammalian liver is the main organ of glutathione synthesis. In the liver of the rats of group 2, the reductive glutathione content exceeded the corresponding indices of the control group by 35%. In the group of older rats the increase amounted to 32%.

Commonly, 80–90% of glutathione is captured and split by the kidneys due to extremely high activity of gamma-glutamyl transpeptidase. Under the influence of Sinestrol, a reductive depletion of the glutathione pool was observed in the kidneys. In the experimental rats (group 2), the reductive glutathione content was reduced by 28% compared with the control.



Table 1. Indices of the brain tissue of rats of different age exposed to synthetic estrogen

Indices	Experiment variants			
	Group 1	Group 2	Group 3	Group 4
Reductive glutathione, mM/g tissue	221 ± 11.05	154.75 ± 7.40*	190 ± 9.50	203.25 ± 10.16
Glutathione reductase, mnM/g tissue	195.2 ± 9.76	168.25 ± 8.41*	174 ± 8.70	200.5 ± 10.03*
Glutathione transferase, mkM/ min. g protein	51.14 ± 2.56	45.77 ± 2.29*	61.06 ± 3.05	57.27 ± 2.86
TBA-active products, nM/g protein	28.61 ± 1.40	38.26 ± 1.90*	24.26 ± 0.12	31.42 ± 1.57*
Total antioxidant activity, standard units	34.35 ± 1.71	21.98 ± 1.10*	27.52 ± 1.38	17.17 ± 0.85*
Superoxide dismutase optic unit/min. g protein	888.46 ± 44.4	764.08 ± 38.2*	543.91 ± 27.2	660.85 ± 33.4*
Glutathione peroxidase, mkM/ min. g protein	26.94 ± 1.347	28.02 ± 1.40	32.75 ± 1.64	38.97 ± 1.95*

\* Significant differences between the indices of control and experimental groups,  $p \leq 0.05$ .

In group 4, the glutathione decline by 17% was recorded. The decrease of glutathione is an indicator of antioxidant system exhaustion.

In the blood serum of group 2 the glutathione content was 77% and in group 4 – 89% of the indices of the corresponding control groups.

In the brain of the control group of rats the glutathione index was 8% higher than the index of the relevant experimental group. The difference between the glutathione content in groups 3 and 4 was 8%. In the brain of rats treated with hormones, the trend of growth of thiol compounds was observed.

In the liver of the experimental 4.5-month-old females, the glutathione reductase activity was 11.5% lower compared to the control. In 7.5-month-old animals, the enzyme activity was 48% higher than in intact females. In the kidneys of females of both experimental groups treated to alimentary estrogen exposure, activity of glutathione reductase did not differ significantly from the control. In the brains of the rats from group 2 inhibition of enzyme activity (13%) was observed. In the brains of older females, the activating enzyme was 2%. In the blood serum of the females from group 2, the glutathione reductase activity was 83%, and in group 4 – 92% of the reference index.

Glutathione transferases play the primary role in the neutralization of secondary peroxidation products and other oxidized compounds. They are conjugated with glutathione basic and most toxic products of lipid peroxidation. These recovery compounds carried out by glutathione transferase prevent progression of peroxidation and the emergence of secondary metabolites. In the liver of 4.5-month-old and 7.5-month-old experimental animals the enzyme activity increased by 18% and 21%, respectively.

There was a trend to activate enzymes in the kidney by 9% (group 2) and by 5% (group 4). The glutathione transferase activity was suppressed in the brain of younger animals by 10.5% and older ones by 6%. In group 2, the activity of the serum enzyme was inhibited by 20%, in group 4 by 8%.

SOD inactivates oxygen radicals occurring in biological reactions of the electron transfer, xenobiotics influence, or various diseases.

In experimental groups of pubertal and mature rats the trend of the increase in the SOD activity by 9% and 5%, respectively, was determined in the liver. In the kidneys of pubertal rats the superoxide dismutase activity grew by 5%. In the experimental group 4, enzyme activation was 17%. In the brains of the group 2 rats, enzyme



inhibition of 14% was observed while in the animals of group 4 the SOD activity increased by 21.5%. The following results were obtained for blood serum: in younger female rats the enzyme activity increased by 32%, in older ones by 15%.

Reducing lipid peroxides in the corresponding alcohols and carrying out the splitting of hydrogen peroxide with the subsequent formation of water, glutathione peroxidase protects the body from oxidative damage.

In determining glutathione peroxidase using glutathione as a donor of H<sup>+</sup>, the activity decreased by 5% in experimental group 2 compared to the control index. In the liver of mature animals, no probable deviations from control were found. In the kidneys, the activity of enzyme inhibition was fixed and its extent was dependent on the age of the animals: in the experimental group of 4.5-month-old rats it was 22%, in 7.5-month-old ones 14% of the intact groups. In the serum of groups 2 and 4, the trend of activity increase by 12% and 5%, respectively, was determined.

The study showed weak activation of glutathione peroxidase (by 4%) in pubertal rats and an increase in the activity of the enzyme by 19% in the brain of mature females.

An investigation into the integral indicator of the antioxidant status – total antioxidant activity in the organs of female rats – indicated a different potential of protection systems depending on age. Thus, in kidneys, blood serum, and the liver an increase in the antioxidant system activity was marked: 4.5-month-old females by 4%, 21% and 18%, 7.5-month-old animals by 30%, 29% and 14%, compared to control groups. At the same time, the study of the brain showed quite a substantial inactivation of antioxidant index of 36% in the pubertal and of 38% in mature females.

In the AChE study, its activity was higher by 22% in the brains of younger rats compared to the control index. In the older group of experimental females, the enzyme activation by 15% was observed. When comparing enzyme index of 3-month- and 6-month-old animals, the activity increase by 5.3% was observed (Table 2).

In blood serum and liver tissue, the inhibition of enzyme activity was determined. In pubertal animals the indicator was significantly different from the corresponding control. For mature females the deviation of activity was less pronounced.

## DISCUSSION

Alimentary estrogen exposure caused an amplification of lipid peroxidation in pubertal animals and mature females. There was a different degree of peroxidation intensification depending on the age and examined organs. The maximum excess of the indices of the intact rats was noted in serum. The kidneys were the most resistant to the impact of exogenous estrogen. In the brains and liver of younger females the reaction of the oxidative system exceeded the power of the response in the organs of sexually-mature animals.

A marked decrease of restored glutathione in the kidneys of the researched female groups demonstrates the risk of violations of the detoxification system. This can lead to an accumulation of free radicals that may trigger factors of various diseases in the future.

Age-dependant organ discreteness to changes in the activity of antioxidant enzymes was observed. This indicates an imbalance of enzymes of the glutathione system. Given the involvement of the glutathione system in the estrogen inactivation by conjugation in their

Table 2. **The impact of nutritional effect of estrogen on the activity of acetylcholinesterase**

Investigated organ	Group 1	Group 2	Group 3	Group 4
Blood serum, mM/mg protein/L	4.52 ± 0.31	4.07 ± 0.28*	4.74 ± 0.36	4.40 ± 0.33
Brain, mM/ mg protein/ min	13.35 ± 0.80	16.23 ± 0.92*	14.06 ± 0.95	16.17 ± 1.03*
Liver, mM/ mg protein/ min	2.77 ± 0.21	2.53 ± 0.25*	2.91 ± 0.18	2.73 ± 0.22

\* Significant differences between the indices of control and experimental groups,  $p \leq 0.05$ .

reactions catalyzed by glutathione transferase, reducing its activity can result in the accumulation of intermediate metabolites and in damage to intracellular structures, mainly DNA.

An imbalance of SOD – glutathione peroxidase links – leads to an accumulation of peroxides, a symptom of endogenous intoxication more pronounced in younger females.

Admission of food hormones to the organism-caused changes of the prooxidant-antioxidant system in the brain and may be followed by a lesion of transmission of information signal. In the brain tissue, the reductive glutathione concentration is at a fairly constant rate. Therefore reducing tripeptide in the brain of a pubertal rat is a negative prognostic criterion of destruction of the afferent integrated system. The estrogen effect concerns the entire central nervous system and includes signaling pathways that intersect with other mediators, such as neurotransmitters and neuromodulator, thus affecting the processes associated with autonomic regulation, neuroprotection, and the like.

The accumulation of toxic metabolites can cause damage on both the cellular and organ levels.

The effects of AChE activity reduction by nutritional estrogen were fixed in the serum and liver. That could be caused by deterioration of the synthesizing function of liver, in particular by stagnation in the liver and kidneys. The enzyme activity was lower in pubertal females compared to mature rats, indicating a lower efficiency of neurotransmitter transmission in the corresponding cholinergic neurons in rats from the younger group. In the future, it can result in serious disturbances in the functioning of the body.

Preserving the enzyme activity on the cerebral cortex level proves the stability of this department. Activation of the enzyme in the cerebral cortex is an evidence of high functional plasticity of the brain tissue. At the same time, the tension in the cholinergic neurotransmitter system was higher in younger rats. Marked changes in their physiological importance are compensatory in nature; according to the experiment and to references in the literature

(Mariani et al., 2011) they are excessive for nerve cells of rat brain and result in a restructuring of the nervous tissue. This is a phenomenon that may trigger the emergence of new pathological states.

## CONCLUSIONS

The state of peroxidation processes and AChE activities in females exposed to xenoestrogen, suggests the possibility of the restructuring of the functioning of mediator and enzyme systems, and of additional enhancement of pathological symptoms. It can be assumed that the effects listed above are able to initiate endocrine disorders, simulate reactions dependant on steroid hormone receptors and independent on the receptor of pathological processes. Regulation of different vectors provides for a different impact on the interaction cell-cell and cell-extracellular matrix and soluble factors. In the body, the identified phenomena can cause inhibition of apoptosis as one of the main mechanisms for removing damaged cells from the population. In the future, such phenomena may trigger a reduction of the potential of compensatory mechanisms, including adaptation and apoptosis, affect the regulation of brain homeostasis, and directly influence the intracellular processes during development and cell differentiation, thus compromising the health of the consumers.

Pubertal females were more sensitive to nutritional synthetic estrogen than mature animals, proving that age is another factor in xenoestrogen exposure. Due to the changes in the rates of reactions of detoxification but not of the metabolism of estrogen received into the organism, particularly with food, with age the animals were less susceptible to the effects of these substances.

Received 13 January 2017

Accepted 25 May 2017

## References

1. Arditsoglou A, Voutsas D. Determination of phenolic and steroid endocrine disrupting

- compounds in environmental matrices. *Environ Sci Pollut Res Int.* 2008; 15(3): 228–36.
2. Boberg J, Taxvig C, Christiansen S, et al. Possible endocrine disrupting effects of parabens and their metabolites. *Reprod Toxicol.* 2010; 30: 301–12.
  3. Cassidy A. Potential risks and benefits of phytoestrogen-rich diets. *Int J Vitam Nutr Res.* 2003; 73(2): 120–6.
  4. Courant F, Antignac JP, Laille J, et al. Exposure assessment of prepubertal children to steroid endocrine disruptors. 2. Determination of steroid hormones in milk, egg, and meat samples. *J Agric Food Chem.* 2008; 14, 56(9): 3176–84.
  5. Courant F, Antignac JP, Maume D, et al. Determination of naturally occurring oestrogens and androgens in retail samples of milk and eggs. *Food Addit Contam.* 2007; 24(12): 1358–66.
  6. Darbre PD, Charles AK. Environmental oestrogens and breast cancer: evidence for combined involvement of dietary, household and cosmetic xenoestrogens. *Anticancer Res.* 2010; 30: 815–27.
  7. De Coster S, van Larebeke N. Endocrine-disrupting chemicals: associated disorders and mechanisms of action. *J. Environ. Public Health.* 2012: 696–713.
  8. Dhaini HR, Nassif RM. Exposure assessment of endocrine disruptors in bottled drinking water of Lebanon. *Environ Monit Assess.* 2014; 186(9): 5655–62.
  9. Hercegová A, Dömötöróvá M, Matisová E. Sample preparation methods in the analysis of pesticide residues in baby food with subsequent chromatographic determination. *J Chromatogr A.* 2007; 15, 1153(1–2): 54–73.
  10. Hu J, Du G, Zhang W, et al. Short-term neonatal/prepubertal exposure of dibutyl phthalate (DBP) induced advanced puberty timing and affected hypothalamic kisspeptin/GPR54 expression differently in female rats. *Toxicology.* 2013; 9(20): 244–8.
  11. Hulin M, Bemrah N, Nougadère A, et al. Assessment of infant exposure to food chemicals: the French Total Diet Study design. *Food Additives and Contaminants: Part A.* 2014; 31(7): 1226–39.
  12. Kolf-Clauw M, Ayouni F, Tardieu D, Guerre P. Variations in zearalenone activation in avian food species. *Food Chem Toxicol.* 2008; 46(5): 1467–73.
  13. Kujawa J, Pasternak K, Zavodnik I, et al. The effect of near-infrared MLS laser radiation on cell membrane structure and radical generation. *Lasers Med Sci.* 2014; 29(5): 1663–8.
  14. Kummer S, von Gersdorff G, Kemper MJ, Oh J. The influence of gender and sexual hormones on incidence and outcome of chronic kidney disease. *Pediatr Nephrol.* 2012; 27(8): 1213–9.
  15. Lykholat T, Lykholat O, Antonyuk S. Immunohistochemical and biochemical analysis of mammary gland tumours of different age patients. *Cytol & Genet.* 2016; 50(1): 32–41.
  16. Mariani A, Jeandel C, Paris F, Ecochard R. Puberty and pubertal growth dynamics in children with idiopathic short stature. *J. Pediatr. Endocrinol Metab.* 2011; 24(6): 319–25.
  17. Massart F, Saggese G. Oestrogenic mycotoxin exposures and precocious pubertal development. *Int J Androl.* 2010; 33(2): 369–76.
  18. McEwen BS. Sex, stress and the brain: interactive actions of hormones on the developing and adult brain. *Climacteric.* 2014; 17(2): 18–25.
  19. Meulenber EP. Phenolics: occurrence and immunochemical detection in environment and food. *Molecules.* 2009; 19, 14(1): 439–73.
  20. Omoruyi IM, Kabiersch G, Pohjanvirta R. Commercial processed food may have endocrine-disrupting potential: soy-based ingredients making the difference. *Food Additives and Contaminants: Part A.* 2013; 30(10): 1722–7.
  21. Petrone AB, Simpkins JW, Barr TL. 17 $\beta$ -estradiol and inflammation: implications for ischemic stroke. *Aging Dis.* 2014; 1; 5(5): 340–5.
  22. Rashid H, Ahmad F, Rahman S, et al. Iron deficiency augments bisphenol A-induced oxidative stress in rats. *Toxicology.* 2009; 256: 7–12.



23. Rochester JR. Bisphenol A and human health: a review of the literature. *Reprod. Toxicol.* 2013; 42: 132–55.
24. Shearwood AM, Chia T, Davies SM, et al. Estrogen-mediated regulation of mitochondrial gene expression. *Mol Endocrinol.* 2015; 29(1): 14–27.
25. Tomar RS, Shiao R. Early life and adult exposure to isoflavones and breast cancer risk. *J. Environ. Sci. Health C Environ. Carcinog Ecotoxicol Rev.* 2008; 26(2): 113–73.
26. Wielogórska E, Elliott CT, Danaher M, Connolly L. Validation and application of a reporter gene assay for the determination of estrogenic endocrine disruptor activity in milk. *Food Chem Toxicol.* 2014; 69: 260–66.
27. Wielogórska E, Elliott CT, Danaher M, Connolly L. Endocrine disruptor activity of multiple environmental food chain contaminants. *Toxicol In Vitro.* 2015; 29(1): 211–20.

**Oleh Yermishev, Tatyana Lykholat,  
Olena Lykholat**

**MAISTINIO SINTETINIO ESTROGENO  
POVEIKIS SKIRTINGO AMŽIAUS ŽIURKIŲ  
LĄSTELĖS KOMPENSACINIAM  
MECHANIZMUI**

*Santrauka*

Daugelis teršalų yra susiję su jų galimu estrogeniniu poveikiu, todėl klasifikuojami kaip endokrini-

nės sistemos veiklą trikdantys junginiai, įskaitant daugelį organinių junginių. Estrogenai yra moteriški lytiniai hormonai, reguliuojantys specifines lytines funkcijas. Juos gamina kiaušidžių folikulai, placenta, iš dalies antinksčių žievė ir sėklidės. Ksenoestrogenai yra pramoninės cheminės medžiagos, naudojamos polikarbonatinio plastiko, epoksidinės dervos, maisto pakuočių ir dangų, odontologinių hermetikų, antimikrobinių preparatų, medikamentų, asmens higienos priemonių gamybai. Šio tyrimo metu buvo tiriama skirtingo amžiaus eksperimentinių žiurkių, paveiktų maistiniu sintetiniu estrogenu, lipidų peroksidacija, antioksidacinės sistemos komponentai ir cholinerginė neuromediatorių sistema. Atsižvelgiant į peroksidacijos procesus ir acetilcholinesterazės aktyvumą patelių organizme panaudojus ksenoestrogenus, galima daryti prielaidą dėl galimo mediatoriaus ir fermentinės sistemos restruktūrizuojančio funkcionavimo, papildomai stiprinančio patologinius simptomus. Nustatyta, kad amžius yra vienas iš ksenoestrogenų poveikio veiksnių, nes bręstančios patelės buvo daug jautresnės maistiniam sintetiniam estrogenui nei subrendę gyvūnai. Dėl reakcijų greičio pokyčių, atsiradus detoksikacijai, bet ne metabolizmui, ypač su maistu estrogeną gaunantys gyvūnai su amžiumi buvo mažiau jautrūs minėtam poveikiui.

**Raktažodžiai:** ksenoestrogenas, lipidų peroksidacija, antioksidacinė sistema, cholinerginis reguliavimas, amžius